

Cichlosoma citrinellum (p. 25).

A specimen has been received from Juan Viñas, Costa Rica (*Underwood*). This species is recorded from Lago de Guila and Lago de Asososca in Nicaragua by Meek (Publ. Columbian Mus., Zool. vii. 1907, p. 152).

34 (A). **Cichlosoma granadense**.

Cichlasoma granadense, Meek, Publ. Columbian Mus., Zool. vii. 1907, p. 121.

Apparently intermediate between *C. citrinellum* and *C. erythraeum*, with the lips and dentition of the former and the low spinous dorsal fin of the latter.

Hab. LAKES MANAGUA AND NICARAGUA.

35 (A). **Cichlosoma dorsatum**.

Cichlasoma dorsatum, Meek, Publ. Columbian Mus., Zool. vii. 1907, p. 123.

Differs from *C. erythraeum* in the longer pectoral fin, which is a little longer than the head and extends to the soft-rayed portion of the anal fin.

Hab. NICARAGUA, Lake Managua.

Cichlosoma affine (p. 26).

This species is recorded by Barbour and Cole (Bull. Mus. Comp. Zool. l. 1906, p. 158) from near Progreso in Yucatan.

46 (A). **Cichlosoma centrale**.

Cichlasoma centrale, Meek, Publ. Columbian Mus., Zool. vii. 1907, p. 94.

Apparently very similar to *C. urophthalmus*, but distinguished by the lower spinous dorsal (longest spine less than $\frac{2}{5}$ the length of head).

Hab. GUATEMALA, Caballo Blanco, Pacific slope.

46 (B). **Cichlosoma tenue**.

Cichlasoma tenue, Meek, Publ. Columbian Mus., Zool. vii. 1907, p. 94.

Differs from *C. centrale* in the slender body (depth $2\frac{3}{4}$ in the length) and the more produced vertical and ventral fins.

Hab. MEXICO, Achotal in Vera Cruz.

Cichlosoma trimaculatum (p. 28).

Recorded from Achotal in Vera Cruz by Meek (Proc. Biol. Soc. Washington, xviii. 1905, p. 245).

47 (A). **Cichlosoma mojarra**.

Cichlasoma mojarra, Meek, Publ. Columbian Mus., Zool. v. 1904, p. 217, fig. 71¹; Proc. Biol. Soc. Washington, xviii. 1905, p. 244².

Hab. MEXICO, San Gerónimo in Oaxaca^{1,2}.