

punctato-striatis, striaque 7^a conspicua; coloribus autem corpore supra æneo-vel olivaceo-nigro, nitido, antennis obscure rufis, articulis 1^o-3^m et palpis clare rufis; capite thoraceque ut in *B. vernali*.

Long. 2½ lin.

Hab. PANAMA, Peña Blanca (*Champion*).

Apparently found in company with *B. vernale*. It much resembles the European (and British) *B. affine*, and is barely distinct from it as a species.

Bembidium mexicanum (p. 148).

To the Mexican localities given add:—Northern Sonora (*Morrison*).

Many examples (labelled *Bembidium*, n. sp.) from Morrison, precisely similar to the bluish form of this species from the 'tierra templada' of Mexico.

Bembidium scintillans (p. 150).

To the localities given add:—NORTH AMERICA, Arizona (*Morrison*). — MEXICO, Northern Sonora (*Morrison*).

Morrison's specimens (labelled *Bembidium*, n. sp.) differ from Central and South Mexican ones by the striæ being a little less impressed for a short distance near the apex, and free from punctures for about a fifth of their length from the apex; in typical examples the punctuation of the striæ continues distinctly further towards the apex. The difference is scarcely worthy of a name as a local form.

ANCHONODERUS (p. 152).

1 (A). **Anchonoderus myops**.

Anchonoderus myops, Reiche, Rev. Zool. 1843, p. 39¹.

Var. *A. duplicatus*. Differt a typo solum macula parva flava prope humerum. (An *A. reichei*, Putzeys, Mitth. Münchener ent. Vereins, ii. p. 57?)

Hab. PANAMA, Bugaba (*Champion*).—SOUTH AMERICA, Colombia¹.

A. myops, according to Reiche's diagnosis, differs from *A. binotatus* in its somewhat larger size and its wider thorax ("thorax capite paullo latior"), *A. binotatus* having "thorax capite vix latior." These features distinguish the numerous series taken by Mr. Champion at Bugaba. The additional red spot near the shoulder is not accompanied by any other difference, and I doubt if the variety be *A. reichei*, Putzeys, from Colombia, as he says the thorax is shorter and "plus brusquement rétréci avant les angles de la base." There is no difference in the form of the thorax from *A. myops*.

3 (A). **Anchonoderus infirmus**.

A. subæneo affinis, at minor graciliorque, elytris subtilissime punctulatis. Castaneo-rufus, capite thoraceque obscurioribus ænescentibus, antennarum dimidio basali, partibus oris pedibusque flavo-testaceis; palpis apice gradatim acuminatis; capite politissimo impunctato; oculis parum prominentibus; thorace quam