

2 (A). **Casnonia lineolata.** (Tab. XIII. fig. 23.)

C. cyanescenti (Chaud.) affinis, quoad formam simillima, sed differt *inter alia* elytris utrinque plagis duabus ex lineolis formatis albo-testaceis, prima ante medium extus versus humerum obliquata interstitia 4^m-8^m, secunda transversa ante apicem interstitia 5^m-8^m, occupanti. Nigra, nitida; antennis et pedibus piceo-fuscis, illarum articulis 1^o-4^m fulvis, 8^o-9^m albis, femoribus basi, tibiis annulo lato, tarsisque partim, testaceo-albis; capite post oculos elongato, recte triangulari impunctato; thorace elongato-angusto, versus medium perparum dilatato, supra transversim haud acute striolato; elytris elongato-oblongis, apice extus breviter dentatis, supra utrinque areis duabus depresso-punctato-striatis, striis apud areas depresso-punctatis profundis ibique interstitiis convexis; interstitiis 3^o et 5^o seriatim punctatis.

Long. 4 lin.

Hab. PANAMA, Bugaba, David (*Champion*).

CALOPHÆNA (p. 163).

Calophæna bicincta (p. 163).

To the localities given add:—PANAMA, Bugaba (*Champion*).

GALERITA (p. 164).

1 (A). **Galerita mustelina.**

G. jelskyi (Chaudoir) proxime affinis et similis, sed differt occipite sparsim punctato fere laevi. Elongata, angusta, nigra, opaca (capite thoraceque subnitidis), tarsis rufo-piceis; capite (♂) quam in *G. jelskyi* (♂) (præcipue spatio inter oculos) conspicue latiori, pone oculos vix prominentes curto, rotundato, convexo, fronte ruguloso-punctato, occipite fere laevi nitido; thorace elongato angusto, antice gradatim modice angustato, crebre punctulato; elytris sicut in *G. jelskyi* elongatis, versus basin angustatis, humeris nullis, costis et costulis geminatis modice elevatis.

Long. 9 lin. ♂.

Hab. COSTA RICA (*coll. Bates*).

I have had this species for many years in my collection among a series of *G. jelskyi* from Ecuador. It is apparently closely allied to *G. stenodera* (Chaud.) from South Brazil, but with narrower or less ovate elytra.

1 (B). **Galerita championi.**

G. mustelinæ affinis; differt capite post oculos multo longiori, semiovali. Elongato-angusta, nigra, subopaca; tarsis fulvo-piceis; capite (♂) ovato, oculis parum prominentibus, post oculos minus convexa, passim grosse, sed haud profunde, crebre punctato; thorace adhuc magis elongato; elytris sicut in *G. jelskyi* elongatis, antice gradatim usque ad basin angustatis, humeris nullis, costis et costulis geminatis acutius elevatis.

Long. 10 lin. ♂.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One example only.

6 (A). **Galerita americana.**

Carabus americanus, Linn. Syst. Nat. ii. p. 671.

Galerita geniculata, Dej. Sp. Gén. Col. v. p. 297¹.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).—WEST INDIES¹. SOUTH AMERICA, Cayenne¹.