

DIAPHORUS (p. 166).

1 (A). **Diaphorus elegans.** (Tab. XIII. fig. 21.)

Diaphorus elegans, Chaudoir, Bull. Mosc. 1862, iv. p. 317¹.

Hab. PANAMA, Peña Blanca (*Champion*).—SOUTH AMERICA, Amazons¹, Sta. Catharina, S. Brazil.

BRACHINUS (p. 169).

13 (A). **Brachinus** — ?

Hab. MEXICO, Presidio (*Forrer*).

A large species, which, in the present confused state of the literature relating to the allied North-American species, it is not possible to determine.

14. **Brachinus fidelis.**

Brachinus fidelis, Leconte, Proc. Acad. Phil. 1862, p. 524¹.

Hab. NORTH AMERICA, New Mexico, Arizona, Lower California¹.—MEXICO, Northern Sonora (*Morrison*).

15. **Brachinus hirsutus**, n. sp.

Quoad formam *B. mexicano* simillimus, sed differt corpore supra (præcipue thorace) longe et dense pubescenti. Testaceo-rufus, antennis concoloribus, abdominis margine atro-fusco, elytris olivaceis sutura et costis 1^a-4^m solum distincte et acute elevatis; thorace sicut in *B. mexicano* lato et postice valde constricto sed subopaco.

Long. 4 lin.

Hab. MEXICO, Pinos Altos in Chihuahua (*Hepburn*).

TETRAGONODERUS (p. 171).

Tetragonoderus mexicanus (p. 173).

To the localities given add :—PANAMA, Tolé (*Champion*).

MASOREUS (p. 174).

Masoreus piceolus (p. 175).

To the localities given add :—PANAMA, Tolé (*Champion*).

PHLŒOXENA (p. 176).

Phlœoxena graphiptera (p. 177).

Var. *b.* (*Ph. limbicollis*). Thoracis limbo lato laterali fulvo.

Hab. PANAMA, Bugaba (*Champion*).