

APENES (p. 188).

11. *Apenes calligramma*. (Tab. XIII. fig. 22.)

Parva, gracilis, parum nitida, castaneo-fusca; capite fere nigro; palpis, antennis, pedibus elytrorumque signaturis fulvis; capite ovato, post oculos rotundato-tumido, supra crebre haud profunde striguloso; thorace cordato-quadrato angusto, ante basin sinuato, angulis posticis fere rectis, lobo basali perparum producto, supra subtiliter confuse striguloso vel coriaceo; elytris elongato-oblongis (postice parum dilatatis), apice sinuatis, macula humerali lineolam apud interstitium 6^m emittente, fascia valde arcuata (ab humero usque fere ad medium curvata, deinde per interstitium 2^m usque ad basin recurvata), fasciola maculari ante apicem (interstitia 5^m-8^m occupante), macula apicali marginibusque angustis, fulvis.

Long. 2½ lin.

Hab. PANAMA (*Champion*).

12. *Apenes nebulosa*.

Apenes nebulosa, Leconte, Proc. Acad. Nat. Sci. Phil. 1866, p. 364; Horn, Trans. Amer. Ent. Soc. x. 1882, p. 158¹.

Hab. NORTH AMERICA, Arizona, Lower California¹.—MEXICO, Northern Sonora (*Morrison*).

APRISTUS (p. 191).

Apristus tropicalis (p. 191).

To the localities given add:—PANAMA, Bugaba, Tolé (*Champion*).

Three examples taken by Mr. Champion differ from *A. mexicanus* and *A. longulus* pretty nearly as required by Motschulsky's description, and in accordance with the observations in my remarks on the species. It can, however, scarcely be specifically different from *A. mexicanus*.

PLOCHIONUS (p. 197).

Plochionus pallens (p. 198).

M. Sallé informs me that *P. valens*, Leconte (New Species Col. i. p. 5), from Pennsylvania, is to be added to the synonymy of this species.

ONOTA (p. 198).

2. *Onota fulvella*. (Tab. XIII. fig. 25.)

O. angulicollis affinis, sed differt statura majore, colore toto testaceo-rufo, vitta laterali viridi-ænea excepta, ab humeris usque ad elytrorum dimidium extensa, interstitia 8^m-9^m occupante; capite et thorace sicut in *O. angulicollis*; elytris relative longioribus, elongato-oblongis, striis profundius impressis vix conspicue punctulatis, interstitiis sat convexis.

Long. 3½ lin.

Hab. PANAMA, Bugaba (*Champion*).

PHILOPHEUGA (p. 202).

Philopheuga viridicollis (p. 202).

Var. Violascenti-cyanea.

Hab. MEXICO, Northern Sonora (*Morrison*).

BIOL. CENTR.-AMER., Coleopt., Vol. I. Pt. 1, August 1884.