

ÆGOPSIS (p. 337).

Ægopsis curvicornis (p. 337).

Ægopsis westwoodii, Thoms. Mus. Scient. p. 17, t. 6. fig. 2¹.

M. Sallé informs me that the only other described species of this genus, *Æ. westwoodii*, Thoms., is not from the interior of Brazil as Thomson stated¹ but from Panama. This is confirmed by the fact that the slight differential features mentioned by Thomson are seen in at least one of our Panama examples.

ISCHNOSCELIS (p. 343).

Ischnoscelis dohrni (p. 343).

Mr. Flohr informs me that an example of this rare species has been taken at Tonila.

COTINIS (p. 346).

Cotinis mutabilis (p. 346).

Var. 2. *C. mexicana* (p. 346).

To the Mexican localities given, add:—Omiteme 8000 feet, Amula 6000 feet, Venta de Zopilote 2800 feet, and Xucumanatlan 7000 feet, all in Guerrero (*H. H. Smith*).

All these examples have a pointed clypeal horn.

Var. 8. *C. atrata* (p. 347).

Add to the Mexican localities:—Amula 6000 feet, and Venta de Zopilote 2800 feet, in Guerrero (*H. H. Smith*).

13 (A). **Cotinis salicis**. (Tab. XXIV. fig. 25.)

C. antonii affinis et simillima, sed valde differt thoracis lobo obtuso sat late truncato. Nigra, polita, pubescentia subtus nigra; caput quadratum, fronte late concava, punctata et nigro-setosa, carina centrali vix elevata et obtusa usque ad clypeum continuata ibique gradatim subsidente; clypeo lato, margine antico valde reflexo recto; thorace elytrisque, præcipue lateribus, sparsim punctatis. Pygidium strigulosum, plagis duabus cretaceis. Processus mesosternalis quam in *C. antonii* paullo longior, apice rotundato. ♂.

Long. 20 millim.

Hab. MEXICO, near the city (*Flohr*).

In decayed willow-trees (*Flohr*).

13 (B). **Cotinis rufipennis**.

Cotinis — ? (no. 18), antea p. 353.

C. antonii affinis. Oblongo-quadrata, nigra, velutino-opaca, subtus (et capite) glabra polita, pectore pedibusque nigro-hirsutis, elytris (sutura angusta nigra excepta) miniato-rufis; clypeus sicut in *C. antonii* margine antico recto oblique reflexo, fronte quadrata, concava, sparsim punctata carinaque alta acuta longe ante clypei apicem subabrupte terminata; antennæ nigrae; thorax trapezoideus, sparse punctatus, disco lævi, ante medium paullo dilatatus, lobo basali mediocre, obtuse rotundato scutelloque conspicuo acute triangu-