

few very minute scattered punctures, very narrowly, subangularly extended on either side behind the eyes; antennæ about three-fourths the length of the body, comparatively stout, thickening a little outwardly, joint 2 moniliform, shorter than 1, 3 as stout as 2, and about equal to it in length, 3-10 gradually increasing in length and width, all longer than broad, 11 ovate, acuminate, nearly twice as long as 10; prothorax transverse, moderately convex, the sides parallel behind and rounded and converging in front, the disc with a very deep, sharply defined, transverse groove before the base (extending across four-fifths of the entire width of the prothorax, and abruptly limited on either side), and an equally deep oblique groove on either side anteriorly, the surface with minute scattered punctures; elytra sparsely, moderately finely punctate, without trace of depression on the disc; femora unarmed, the tibiæ almost straight.

Length 1 millim.

*Hab.* MEXICO, Jalapa (*Flohr*).

#### NOTOXUS (p. 203).

##### **Notoxus ventralis** (p. 207).

To the Mexican localities given, add:—Mexico city and Chalpultepec (*Flohr*).

Numerous examples of both sexes have been received from Mr. Flohr. The female has the elytra abruptly sinuate-truncate at the apex, with the sutural angles slightly produced. The male has a deep foveate depression at the apex of the last ventral segment. The arrangement of the elytral pubescence is constant, though the testaceous markings are sometimes indistinct.

#### TOMODERUS (p. 215).

##### **Tomoderus brevicornis** (p. 218).

To the localities given, add:—MEXICO, Yautepec in Morelos (*Flohr*).

##### **Tomoderus brevis** (p. 219).

To the localities given, add:—MEXICO, Motzorongo in Vera Cruz (*Flohr*).

#### ANTHICUS (p. 221).

##### **Anthicus dilatipes** (p. 231).

To the Mexican locality given, add:—Tepetlapa in Guerrero (*Flohr*).

A single male example, agreeing perfectly with the type.

##### 15 (A). **Anthicus sturmi**.

*Anthicus elegans*, La Ferté, Monogr. Anthic. p. 117, t. 27. figg. 12, 14<sup>1</sup>; Lec. Proc. Acad. Phil. vi. p. 96<sup>2</sup> (nec Stev.).

*Anthicus sturmi*, La Ferté, loc. cit. p. 304.

*Hab.* NORTH AMERICA, Carolina<sup>1</sup>, Georgia, and Missouri<sup>2</sup>.—MEXICO, Taxco (*Flohr*).

One female specimen, agreeing very nearly with La Ferté's description.