

**Cholus pilicauda** (p. 303).

To the localities given, add :—COSTA RICA, Caché (*Biolley*), Carrillo (*Underwood*).

Four specimens.

**Cholus leucogaster** (p. 304).

To the locality given, add :—COSTA RICA, Zeut, Atlantic slope, 20 metres (*Pittier*).

Three more specimens of this species have been received, including both sexes. The female has the rostrum much smoother than in the male, and its median carina short.

**Cholus pallidiventris** (p. 305).

To the locality given, add :—COSTA RICA, Carrillo (*Underwood*).

One specimen, differing from those described from Nicaragua in having the scales on the sides of the body beneath ochreous, instead of white.

37 (A). **Cholus chrysalis**. (Tab. XXXV. figg. 28, 28 a.)

*Cholus chrysalis*, Heller, Stett. ent. Zeit. 1906, p. 15, t. 1. fig. 6<sup>1</sup>.

Elongate, narrow, subfusiform, flattened above, moderately shining, piceous or rufo-piceous, the head, antennæ (the club excepted), rostrum, and legs rufescent, the apices of the femora black; sparsely clothed with small ochreous scales, the sides of the body beneath densely covered with coarser whitish scales; the femora and tibiæ shining, almost bare. Head finely punctate, rugulose above the eyes, the latter rather prominent and well-separated; rostrum moderately stout, arcuate, reaching the mesosternum, finely punctate, carinate, and sulcate at the base, and almost smooth thence to the tip, the antennæ inserted at some distance beyond the middle, joint 1 of the funiculus about as long as 2 and 3 united. Prothorax a little broader than long, somewhat rounded at the sides, narrowing from near the base, feebly constricted in front; closely, transversely granulate. Scutellum about as broad as long. Elytra elongate-cordate, not wider than the prothorax, conjointly rounded at the apex; closely granulate and indistinctly seriate-punctate, the outer interstices narrowly costate. Mesosternum flattened between the coxæ. Metasternum and abdomen with a broad, almost bare, smooth space down the middle, the first ventral segment broadly depressed along the centre. Legs long and slender, comparatively smooth; femora each with an acute tooth, the hind pair reaching a little beyond the apex of the elytra; tibiæ armed with a short claw at both the outer and inner apical angles, the anterior pair dilated at about the middle within.

Length 9–10 $\frac{1}{8}$ , breadth 3–3 $\frac{1}{2}$  millim. ( $\sigma$ ?)

*Hab.* COSTA RICA, San Carlos (*Mus. Dresden*<sup>1</sup>), Zeut, Atlantic slope, 20 metres (*Pittier*).

We have received three specimens of this species. *C. chrysalis* somewhat resembles *Erethistes zygopoides*; it is, however, much smaller, and has the legs shorter, smoother, and much more slender, the tibiæ unguiculate, the sides of the body beneath densely clothed with whitish scales, &c. The vestiture of the under surface is similar to that of *Cholus pallidiventris*.

## ERETHISTES (p. 310).

**Erethistes cyanipes** (p. 311).

To the localities given, add :—COSTA RICA, San José 1160 metres (*Biolley*).