

Fam. HISPIDÆ.

A P P E N D I X*.

SINCE the completion of the Supplement to this Family in May 1886 by the late J. S. Baly many specimens of Hispidæ have been received from Southern Mexico, and we take this opportunity, before closing the volume, to include here this additional material. The extensive collection made by Mr. H. H. Smith, principally in the States of Guerrero, Vera Cruz, and Tabasco, includes several new species, and adds much to our knowledge of the geographical distribution of many of these interesting insects. Mr. Gaumer has also sent us two novelties from Yucatan.

STENISPA (p. 4).

Stenispa sallæi (p. 4).

To the Mexican localities given, add:—R. Papagaio in Guerrero, Teapa in Tabasco (*H. H. Smith*).

Many specimens of both sexes were obtained by Mr. Smith at Teapa.

CEPHALOLEIA (pp. 8, 120).

Cephaloleia fulvolimbata (p. 24).

To the localities given, add:—MEXICO, Teapa in Tabasco (*H. H. Smith*).

Numerous specimens of the typical form.

CEPHALODONTA (pp. 32, 120).

Cephalodonta erudita (p. 39).

To the localities given, add:—MEXICO, Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*).

Two specimens of the var. A, hitherto only known from Nicaragua.

CHARISTENA (p. 45).

Charistena championi (p. 46).

To the localities given, add:—MEXICO, Teapa in Tabasco (*H. H. Smith*).

* By G. C. CHAMPION.