

Chalepus roseus (p. 58).

To the Mexican localities given, add:—Omilteme and Amula in Guerrero (*H. H. Smith*).

Chalepus fryi (p. 59).

To the localities given, add:—MEXICO, Omilteme and Chilpancingo in Guerrero, Atoyac in Vera Cruz (*H. H. Smith*).

This insect is probably a variety of *C. roseus*.

19 (A). **Chalepus flavipes**. (Tab. XIII. fig. 24.)

Moderately elongate, cuneiform, slightly shining; above and beneath rufo-fulvous, the elytra with an indeterminate transverse fascia before the apex dilute sanguineous, the lateral margin (except at the apex), and also the apical margin, flavous, the eyes and antennæ black; the legs flavous, the tarsi fuscous. Head opaque, with a few scattered punctures; the front moderately produced, the interocular space longitudinally sulcate in the middle. Antennæ short, not one third the length of the body, thickening outwardly; the joints cylindrical, 3 slightly longer than 2 or 4, 6–10 transverse, 11 longer than 10, acuminate. Prothorax transverse, convex; the sides converging from the base, the anterior angles rather obtuse; the surface closely covered with coarse, foveolate punctures, separated by fine raised reticulations. Elytra moderately long, gradually widening from the base to the rounded posterior angle; the apices conjointly, obtusely rounded; the lateral and apical margins finely serrulate; each elytron with eight, at the base and apex with nine, rows of punctures; the second, fourth, and sixth interspaces, together with the suture, costate, the second more broadly and strongly than the others. Legs rather short.

Length $2\frac{1}{4}$ lin.

Hab. MEXICO, Cuernavaca in Morelos (*H. H. Smith*).

One specimen. This insect closely resembles *C. instabilis*. Baly, var. C, but may easily be known from it by the short antennæ and pale legs. The penultimate joints of the antennæ are freely articulated and strongly transverse.

Chalepus marmoratus (p. 60).

To the localities given, add:—Cuernavaca in Morelos, Atoyac in Vera Cruz (*H. H. Smith*).

Chalepus digressus (p. 66).

To the Mexican locality given, add:—Teapa in Tabasco (*H. H. Smith*).

One specimen, differing from the type in having the apical black patch of the elytra more extended.

Chalepus bellulus (pp. 68, 121).

To the Mexican localities given, add:—Venta de Pelegrino, Tierra Colorada, Tepetlapa, Venta de Zopilote, and Dos Arroyos in Guerrero, Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*).

Chalepus rufithorax (p. 70).

To the Mexican locality given, add:—Tierra Colorada in Guerrero (*H. H. Smith*).

One specimen, agreeing perfectly with the type.