

**Eudamus albofasciatus** (II. p. 280).

To the localities given, add :—HONDURAS, San Pedro Sula (*Wittkugel*).

**Eudamus alcæus** (II. p. 281).

*Thymele montezuma*, Scudd. Rep. Peabody Ac. Sci. iv. p. 70 (1872) <sup>4</sup>.

To the localities given, add :—MEXICO, Tehuantepec <sup>4</sup>; HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*).

The name *E. alcæus*, Hew., has five years' priority over that of Scudder (cf. *anteà*, p. 636).

17 (A). **Eudamus cachinnans**, sp. n. (Tab. CXII. figg. 3, 4, ♂.)

Alis fuscis, anticis fascia lata mediana per cellulam venis divisa, altera angustiore subapicali curvata, maculis duabus subquadratis inter eas, proxima majore, omnibus hyalinis; posticis striga mediana elongata indistincta grisea, margine interno dense pilis ferrugineis vestitis, ciliis (nisi ad apicem) albis: subtus anticis ut supra, posticis striga elongata alba cæruleo tineta, punctis nigricantibus duobus includente; capite et thorace supra indistincte viridi suffusis.

*Hab.* PANAMA, Chiriqui (*Trötsch, ex Staudinger*).

We have received a male and female of this species from Dr. Staudinger, who sent them under the name of *E. ridens*, Hew., both occurring in Chiriqui. It differs, however, from that insect, of which we now possess both sexes, in having the hyaline spot on the disc of the primaries much less angulate, forming a more regular fascia; the secondaries have the whitish streak on the disc but very faintly indicated, the white tuft of hairs towards the anal angle absent, and the longitudinal band on the underside somewhat broader, and the head and thorax are suffused with green.

19 (A). **Eudamus biolleyi**. (Tab. CXII. figg. 5, 6, ♂.)

*Eudamus biolleyi*, Mab. Bull. Soc. Ent. Fr. 1900, p. 230 <sup>1</sup>; Godm. & Salv. *anteà*, ii. p. 635 <sup>2</sup>.

Alis nigro-fuscis viridi-suffusis, anticis fascia mediana interrupta per cellulam, altera subapicali angustiore, maculis duabus inter eas, proxima majore, omnibus hyalinis; posticis ciliis (nisi ad apicem) et cauda albis: subtus ut supra, anticis litura cærulea angulum analem versus; posticis striga lata elongata marginem externum versus cæruleo-alba maculis duabus nigris includente.

*Hab.* COSTA RICA <sup>2</sup>, Pecara (*coll. Joannis* <sup>1</sup>).

The Abbé Joannis has been kind enough to lend us the type of this insect, which, as Mabile notes, is a very close ally of the Guatemalan *E. crison*, described and figured in this work from an imperfect female. It differs, however, from that species in having much smaller hyaline spots on the primaries, the median band being narrower and interrupted, the secondaries are without spots on the upperside, the long stripe beneath is nearer the outer margin and somewhat differently shaped, and the tails and part of the fringe are white. It is not improbable that this last-named character will also be found in *E. crison* when perfect specimens are obtained.