

PACHES (II. p. 398).

Paches polla (II. p. 399).

♀ mari similis, sed colore cyanea absente, punctis duobus minutis, uno inter ramos medianos primum et secundum, secundo margini interiori proximo, hyalinis.

To the localities given, add :—GUATEMALA, San Gerónimo (*Champion*).

We now give a description of the female of this species, of which we possess three examples, two from Costa Rica and one from Guatemala.

EANTIS (II. p. 404).

Eantis busirus (II. p. 404).

To the localities given, add :—HONDURAS, San Pedro Sula (*Wittkugel*) ; NICARAGUA, Matagalpa (*Richardson*).

Eantis thraso (II. p. 405).

To the localities given, add :—HONDURAS, San Pedro Sula (*Wittkugel*).

DOBERES (II. p. 407).

1. Doberes hewitsonius.

Achlyodes hewitsonius, Reak. Proc. Ac. Phil. 1866, p. 340 (Nov.)¹.

Doberes mexicanus, anteà, ii. p. 407, t. 87. f. 11².

To the localities given, add :—MEXICO, near Vera Cruz (*Edwards*¹).

Reakirt's description of *A. hewitsonius*¹ was overlooked by us in our first enumeration of the species of Hesperiinae; it antedates that of Felder by three months (cf. *anteà*, p. 635).

EBRIETAS (II. p. 418).

Ebrietas osyris (II. p. 419).

To the localities given, add :—HONDURAS, San Pedro Sula (*Wittkugel*).

THEAGENES (II. p. 428).

In our first enumeration of the species of this genus, two were accidentally omitted: they are described on pp. 459, 460.

STAPHYLUS (II. p. 429).

5. Staphylus mazans.

Carcharodus mazans, Reak. Proc. Ac. Phil. 1866, p. 335¹.

Staphylus ascalaphus, anteà, ii. p. 432, t. 89. ff. 12–15².

Reakirt's types of *C. mazans* were from Vera Cruz, whence we have already recorded this insect (cf. *anteà*, p. 635).