

12. *Coronidia ribbei*, sp. n. (Tab. XLI. fig. 14, ♀.)

Female. Primaries dark brown, crossed from the costal to the inner margin by many waved brown lines, a wide pale yellowish-brown band crossing the wing from the costal to the inner margin close to the anal angle, the outer margin broadly bordered with pale brown, the latter slightly dentated near the apex; secondaries dark brown, with a wide pale yellow band extending from the costal margin near the apex to the inner margin above the anal angle, the outer margin deeply dentated; the underside pale yellow, the primaries with a rather large black spot at the end of the cell, and crossed from the costal to the inner margin by two wide waved black bands, the secondaries crossed only by a central black band, the outer band being broken up into very fine lines: head, thorax, abdomen, antennæ, and legs brown; the palpi brown, with the third joint black. Expanse 2 inches.

Hab. PANAMA, Chiriqui (*Ribbe, in mus. Staudinger*).

A single specimen only of this very distinct species has been communicated by Dr. Staudinger; its nearest ally appears to be *C. genevana*, Westw.

13. *Coronidia ocyclus*.

Coronis ocyclus, Boisd. Lép. Guat. p. 76¹.

Coronidia ocyclus, Westw. Trans. Zool. Soc. x. p. 539².

Coronis dutreuxi, Deyr. Rev. et Mag. Zool. 1874, t. 8. f. 3³.

Hab. ? MEXICO^{1 2}; GUATEMALA^{1 2}; COSTA RICA³.

We are indebted to Mons. C. Oberthür for kindly lending us the type of *C. ocyclus*, Boisd.; it proves to belong to the same species as *C. dutreuxi*, Deyr.

Sect. 4. *Alis posticis haud fasciatis*, Westw.

14. *Coronidia leachii*.

Agarista leachii, Godt. Encycl. Méth. ix. p. 803¹.

Coronis leachii, Guérin, Icon. Règne Anim., Ins. p. 493, t. 83. f. 3²; Gray, in Griffith's Anim.

Kingd., Ins. p. 596, t. 53. f. 3³; Boisd. Sp. gén. des Lép. i. t. 14 (10 B), f. 2⁴; Walk. Cat.

i. p. 39⁵; Guen. Sp. gén. des Lép. ix. p. 22, Uran. t. 1. f. 2 (♂)⁶.

Coronidia leachii, Westw. Trans. Zool. Soc. x. p. 540, t. 88. f. 11⁷.

Hab. MEXICO (*Hartweg*^{5 7}); GUATEMALA, Coban in Vera Paz (*Conradt*), San Gerónimo (*Champion*); PANAMA, Chiriqui (*Ribbe*), Volcan de Chiriqui 3000 to 4000 feet (*Champion*).—BRAZIL^{1 2 6 7}.

Mr. Champion captured a good series of specimens of this insect at San Gerónimo. It varies considerably in size, one of the examples from Chiriqui being hardly more than half the size of some of those from Guatemala. From Guatemala I have both sexes before me; they clearly prove that Guenée was wrong in supposing that *C. leachii* was the male of the yellow-banded *C. jayet*, the markings of the wings scarcely differing in the two sexes.