

## ODOPŒA.

*Odopœa*, Stål, Ann. Soc. Ent. Fr. sér. 4, vol. i. p. 616 (1861); Hem. Afr. iv. p. 1 (1866).

This genus differs from the preceding in having the ulnar veins clearly separated at the base. Stål also adds, "marginibus lateralibus areæ ulnaris interioris parallelis." *Odopœa*, though confined to the Neotropical Region, has a wider area of dispersal than that of *Zammara*. Though not recorded north of Mexico, it extends as far south as the Argentine Republic, and is well represented in the Antilles. About thirteen species are known, six of which have been found in Central America.

1. ***Odopœa montezuma***. (Tab. III. figg. 5, 5 a, 5 b.)

*Zammara montezuma*, Walk. List Hom. i. p. 36. 6<sup>1</sup>.

*Odopœa montezuma*, Stål, Ann. Soc. Ent. Fr. sér. 4, i. p. 617; Öfv. Vet.-Ak. Förh. 1862, p. 484; Stett. ent. Zeit. xxv. p. 58. 383<sup>2</sup>.

*Hab.* MEXICO<sup>1 2</sup>.

The figure is taken from a typical specimen in the collection of the British Museum. We have not yet received it in any of the collections sent to us from Central America.

2. ***Odopœa imbellis***. (Tab. I. figg. 3, 3 a, 3<sup>a</sup>b.)

*Zammara imbellis*, Walk. List Hom. Suppl. p. 2<sup>1</sup>.

*Odopœa imbellis*, Stål, Ann. Soc. Ent. Fr. sér. 4, i. p. 617; Öfv. Vet.-Ak. Förh. 1862, p. 484; Stett. ent. Zeit. xxv. p. 58. 384<sup>2</sup>.

*Hab.* MEXICO<sup>1 2</sup> (*Heller, Mus. Vind. Cæs., Mus. Holm.*); GUATEMALA, Panama (*Champion*).

A Guatemalan specimen is figured.

3. ***Odopœa azteca***, n. sp. (Tab. I. figg. 4, 4 a, 4 b.)

Body pale green. Head with the anterior margin, outer margins of ocelli, and an oblique narrow fascia from outer margins of ocelli to frontal margin of eyes black; ocelli red, eyes ochraceous. Pronotum with a central transverse fascia near anterior margin, two central longitudinal narrow fasciæ on disk, two oblique narrow fasciæ on each side, commencing from behind eyes, and outer and inner borders of dilated lateral margins black. Mesonotum with two large central obconical and curved spots on anterior margin, followed by a central longitudinal fascia on disk, on each side of which is a rounded spot, and on outer side of these central markings four elongate spots, placed in pairs obliquely, black. Abdomen above with a small black central transverse spot on anterior margin of first segment, and four small rounded fuscous spots near lateral margins of third, fourth, fifth, and sixth segments. Lateral margins of apical half of face, apex of rostrum, bases and apices of tibiæ (narrowly), anterior tarsi, excepting base of apical joint, and apices of posterior tarsi black. Tegmina pale hyaline, with the costa and radial veins greenish; basal half of tegmina with the veins greenish, outer half with the veins fuscous; claval margins testaceous, posterior edged with black near base. Wings pale hyaline, veins coloured as in tegmina, but claval margin darker.

Face somewhat depressed, lateral margins transversely striated and centrally and longitudinally sulcated. Lateral margins of pronotum much amplified and obtusely angulated, posterior margins transversely striated. Tympana prominent, subconical, and subacutely angulated anteriorly. Opercula truncate