

#### 47. *Gypona notula*, sp. n.

*Parva*, testacea ; capite lato, oculis magnis, ocellis inter se quam ab oculis longius distantibus, haud producto vel attenuato ; pronoto antice punctis quibusdam nigris instructo, sat longo ; scutello magno, testaceo ; tegminibus haud irroratis, sed sparsim fusco vel nigro notatis ; corpore subtus pedibusque dilute testaceis.

*Mas* segmento ultimo ventrali longo ad apicem rotundato.

*Femina* segmento ultimo ventrali leviter sinuato.

A small testaceous species, with the head broad and not produced, and with bright red ocelli situated nearer to the eyes than to one another ; pronotum rather long, with from four to eight small black spots in front ; tegmina with irregular fuscous or black markings.

*Male* with the last ventral segment long, rounded at the apex.

*Female* with the last ventral segment slightly sinuate.

Long. 6 millim. ; lat. ad hum. 2 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

Two specimens, male and female. Not unlike *G. hieroglyphica*, but smaller and with the ocelli more widely separated.

#### 48. *Gypona bisignata*, sp. n. (Tab. XXI. figg. 23, 23 a, b, ♀.)

Dilute testacea ; capite haud attenuato, ocellis inter se quam ab oculis paulo longius distantibus ; pronoto antice nigro-testaceo ; scutello fusco variegato ; tegminibus nigro sparsim irroratis, maculâ magnâ nigrâ utrinque vix pone medium positâ ; pedibus testaceis.

*Mas* ignotus. *Femina* segmento ultimo ventrali leviter sinuato, fere recto.

Somewhat variable in size ; of a light testaceous colour ; head comparatively narrow, the ocelli large, situated somewhat nearer to the anterior than to the posterior margins, and a little further separated from one another than from the eyes ; pronotum with several dark spots and markings in front, strongly striate transversely ; scutellum variegated with fuscous ; tegmina with small, scattered, dark spots, which are thicker at the suture, and with a large, dark, irregular patch on each just behind the middle ; legs and underside light testaceous, the sides of the chest dark.

*Male* unknown. *Female* with the apical margin of the last ventral segment almost straight, feebly sinuate and very slightly emarginate in the middle.

Long. 8-9 millim. ; lat. ad hum. 2½-3 millim.

*Hab.* MEXICO, Atoyac in Vera Cruz (*Schumann*) ; GUATEMALA, San Gerónimo (*Champion*).

The Guatemalan insect is figured.

#### 49. *Gypona reservanda*, sp. n. (Tab. XXI. figg. 24, 24 a, b, ♀.)

Fusco-testacea ; capite sat longo, sed haud subtus attenuato, ocellis inter se quam ab oculis paullo magis distantibus ; pronoto sat fortiter transversim striato, antice nigro notato ; tegminibus maculis quibusdam parvis nigris et notâ utrinque majore circulari ad marginem pone medium instructis ; corpore subtus pedibusque fuscis.

*Mas* ignotus. *Femina* segmento ultimo ventrali late sed haud profunde emarginato, sinuato, ad medium leviter exciso.

Fusco-testaceous, with the head somewhat produced and pointed, but not attenuated beneath ; ocelli a little further from one another than from the eyes ; pronotum with black markings in front ; tegmina with a few small dark markings and a rather large semicircular spot on each near the margins behind the middle, the veins not marked ; wings fuscous ; underside and legs fuscous.