

Male unknown. *Female* with the apex of the last ventral segment broadly and shallowly emarginate, and sinuate and slightly excised in the centre.

Long. 7 millim.; lat. ad hum. 2 millim.

Hab. MEXICO, Venta de Zopilote in Guerrero 2800 feet (*H. H. Smith*).

50. **Gypona bohemani.**

Gypona bohemani, Stål, Stett. ent. Zeit. xxv. p. 81¹; Spångb. Bihang till K. Sv. Vet.-Ak. Handl. v. 3, p. 52².

Hab. MEXICO, Vera Cruz^{1 2} (*Mus. Holm.*).—COLOMBIA, Bogota².

I have not seen an example of this insect, and the female is apparently unknown.

51. **Gypona præterita**, sp. n. (Tab. XXI. figg. 25, 25 a, b, ♀.)

Brunneo-testaceo, capite haud producto, obtuso, ocellis inter se quam ab oculis longius remotis; pronoto lateribus antice sensim rotundatis, fortiter transversim striatis; scutello magno, postice acuminato; tegminibus punctis quibusdam nigris minutis; corpore subtus pedibusque plus minusve fuscatis.

Mas segmento ultimo ventrali brevi fere recto, valvulis brevibus.

Femina segmento ultimo ventrali late et haud profunde emarginato, levissime sinuato.

Brownish-testaceous, with the head not produced and blunt, the ocelli situated nearer to the front than to the basal margin, more distant from one another than from the eyes; pronotum with the sides rounded anteriorly; tegmina with a few minute, irregular, black markings; underside and legs mostly fuscous.

Male with the last abdominal segment and the genital valves short.

Female with the apex of the last ventral segment of the abdomen broadly and shallowly emarginate and very slightly sinuate.

Long. 8 millim.; lat. 2½ millim.

Hab. MEXICO, Atoyac in Vera Cruz (*H. H. Smith*).

In the collection before me there is also a specimen from Teapa which might perhaps be referred to this species.

52. **Gypona extranea**, sp. n.

Castaneo-brunnea, capite brevi, antice obtuso haud attenuato, ocellis inter se quam ab oculis multo magis distantibus; pronoto unicolori, fortiter transversim striato; tegminibus notis quibusdam nigris vix conspicuis, venis ad apicem evidentioribus.

Mas segmento ultimo ventrali penultimo longiori, ad marginem leviter sinuato, in medio denticulato producto.

Femina segmento ultimo ventrali late leviter emarginato.

Short and rather broad, of a castaneous-brown colour; head short, blunt, and not produced, with the ocelli much nearer to the eyes than to one another; pronotum concolorous, strongly striate transversely; tegmina with a few very small and inconspicuous black markings, and the veins marked at the apex; underside and legs fuscous or fusco-testaceous.

Male with the last ventral segment of the abdomen longer than the penultimate, slightly sinuate at the apex, and produced into a small tooth in the centre.

Female with the last ventral segment broadly and shallowly emarginate.

Long. 8-9 millim.; lat. ad hum. 3 millim.

Hab. GUATEMALA, Volcan de Atitlan (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).