

53. *Gypona exornata*, sp. n. (Tab. XXI. figg. 26, 26 *a*, *b*, ♀.)

Sat robusta, parallela, lète castaneo-brunnea, testaceo et rufo-testaceo variegata; capite brevi, haud producto, antice rotundato, ocellis approximatis, oculis magnis; pronoto testaceo rufo-testaceo variegato, postice brunneo; scutello brunneo, postice albido; tegminibus maculis duabus magnis ad marginem utrinque testaceis, et quibusdam minoribus prope suturam et ad humeros instructis; pedibus rufo-testaceis.

Mas ignotus. *Femina* segmento ultimo ventrali late sed haud profunde sinuatim emarginato.

A somewhat stout, subparallel-sided species, of a bright chestnut-brown colour of different shades; head short, not produced, rounded in front, testaceous, variegated with orange; ocelli approximate; pronotum testaceous, with bright orange markings, and with the hind margin dark brown; scutellum brown, with the hinder part clear yellowish-white; tegmina almost parallel, somewhat rounded at the sides, broadest just behind the middle, with two large testaceous spots on each on the margins, and several small ones near the suture and at the shoulders; legs rufo-testaceous.

Male unknown. *Female* with the last ventral segment broadly emarginate, the emargination being sinuate and not deep.

Long. 7 millim.; lat. ad hum. 3 millim.

Hab. PANAMA, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).

54. *Gypona wallengreni*. (Tab. XXI. fig. 27, ♂.)

Gypona wallengreni, Stål, Stett. ent. Zeit. xxv. p. 82¹; Spångb. Bihang till K. Sv. Vet.-Ak. Handl. v. 3, p. 51².

Hab. MEXICO² (*Mus. Holm.*¹); PANAMA, Volcan de Chiriqui (*Champion*).

Easily known by its very broad, short head and by the wide testaceous border of the dark ferruginous tegmina.

55. *Gypona nana*, sp. n. (Tab. XXI. figg. 28, 28 *a*, *b*, ♀.)

Parva, nitida, castaneo-brunnea, interdum colore obscuriori variegata; capite lato, antice obtuse producto, ocellis inter se quam ab oculis paullo magis distantibus; pronoto sat magno, disco unicolori, vel antice nigro notato, marginibus lateralibus anguste albidis, ad medium transversim striato; scutello magno, postice producto, acuminato; tegminibus margine exteriori albido limbato, limbo nigro-variegato; corpore subtus pedibusque dilute testaceis.

Mas ignotus. *Femina* segmento ultimo ventrali fortiter sinuato ad medium triangulariter exciso.

A very small, shiny, castaneous-brown species, which may at once be known by its size and by the white borders of the pronotum and tegmina, the latter being variegated with black; head broad, the ocelli situated nearer the anterior than the posterior margin, a little more distant from one another than from the eyes; pronotum large, unicolorous, or with dark markings in front; underside and legs light testaceous.

Male unknown. *Female* with the apex of the last ventral segment of the abdomen strongly sinuate and triangularly excised in the centre.

Long. 5 millim.; lat. ad hum. 2 millim.

Hab. MEXICO, Teapa in Tabasco (*H. H. Smith*); PANAMA, Bugaba (*Champion*).

We figure a specimen from Teapa.