

published six years afterwards (1864), he describes two species of the genus from Mexico under the generic name of *Stragania*. We do not possess an example of either of these in our collection, and I include them here under the Gyponidæ on Stål's authority and append his descriptions. They are small insects, with hyaline or subhyaline tegmina, and resemble Jassidæ rather than Gyponidæ.

1. *Stragania humilis.*

Stragania humilis, Stål, Stett. ent. Zeit. xxv. p. 85¹.

"Griseo-albida; fronte, loris, medio verticis thoracisque, scutello, apice excepto, macula magna mesostethii basique coxarum posticarum nigris; tegminibus breviter fusco-pilosus; capite thorace vix angustiore. ♀. Long. 4 millim."

Hab. MEXICO (*coll. Signoret*¹, *Mus. Vind. Cæs.*).

2. *Stragania misella.*

Stragania misella, Stål, Stett. ent. Zeit. xxv. p. 85¹.

"Flavo-testacea, subtus cum pedibus stramineo-albida; tegminibus sordide hyalinis, fusco-pilosulis, scutello, clavo, apice excepto, maculaque subapicali dorsi abdominis nigris; scutelli maculis discoidalibus flavo-testaceis; capite thoracis parte posteriore paullo angustiore. ♂. Long. 4 millim."

Hab. MEXICO, Vera Cruz (*Mus. Holm.*¹).

NOTE.—Since the publication of the preceding page, I find that Stål subsequently included *Stragania* under the Jassidæ as being synonymous with *Macropsis*, Lewis [Trans. Ent. Soc. i. p. 49 (1834)]; the alteration appears in very small type in his 'Hemiptera Africana,' iii. p. 126.