SUPPLEMENT.

THE following species have been described or recorded from Central America since the publication of the preceding pages.

MEMBRACIDÆ.

1. Membracis nebulosa.

Membracis nebulosa, Buckton, Mon. Membr. p. 36, t. 2. fig. 8 (1901) 1.

Hab. Costa Rica 1.

This species seems somewhat doubtful, and Mr. Buckton, in his Monograph, marks it with a note of interrogation. The specimen on which it is described is evidently imperfect, and it is probably one of the many varieties of *M. mexicana*, Guér.

2. Membracis (Phyllotropis) dorsata.

Membracis dorsata, Fabr. Syst. Rhyng. p. 11 (1803) 1. Phyllotropis dorsata, Buckton, Mon. Membr. p. 40 (1901) 2.

Hab. CENTRAL AMERICA 2.—SOUTH AMERICA 1.

According to Buckton, there are several specimens of this insect from Central America in the National Collection; but I think there must be some mistake about this locality, as I have carefully examined all the examples in the Museum collection. Phyllotropis is, at most, a subgenus of Membracis. The same author (l. c. pp. 35, 41) places M. mexicana under both Membracis and Phyllotropis; on page 40 he gives my Membracis humilis as a synonym of Phyllotropis trifasciata, Stål, but on page 43 he describes the species as Phyllotropis humilis, and copies the figure from the 'Biologia' (t. 1. fig. 6) on plate iv. fig. 3.

3. Euchenopa tesselata.

Euchenopa tesselata (tessilata in error), Buckton, Mon. Membr. p. 49, t. 6. fig. 1 (1901) 1.

Hab. Mexico 1.—South America, Rio Janeiro 1.

Described from broken specimens.

4. Sphongophorus inelegans.

Sphongophorus inelegans, Buckton, Mon. Membr. p. 82, t. 15. fig. 6 (1901) 1.

Hab. Mexico 1 (Hope coll., Oxford).