

for description; as, however, the type is not among the insects returned to me after Mr. Buckton's death, I cannot say anything further about it. I have some recollection of noticing the peculiar venation of the tegmina.

13. **Darnoides flavescens.**

*Darnoides flavescens*, Baker, Canad. Ent. xxxix. p. 117 (1907)<sup>1</sup>.

*Hab.* BRITISH HONDURAS, Belize<sup>1</sup>.

14. **Heteronotus vulnerans.**

*Combophora vulnerans*, Germ. Rev. Silb. i. p. 228 (1833)<sup>1</sup>.

*Heteronotus vulnerans*, Buckton, Mon. Membr. p. 141, t. 30. fig. 1 (1902)<sup>2</sup>.

*Hab.* MEXICO, Vera Cruz<sup>2</sup>; GUATEMALA<sup>2</sup>; PANAMA<sup>2</sup>.—BRAZIL<sup>1 2</sup>.

15. **Poppea succinea.**

*Poppea succinea*, Buckton, Trans. Linn. Soc., Zool. (2) ix. p. 331, t. 21. figg. 7, 7 a (1905)<sup>1</sup>.

*Hab.* MEXICO<sup>1</sup> (*Mus. Madrid*).

16. **Ceresa nitens.**

*Ceresa nitens*, Buckton, loc. cit. p. 332, t. 21. figg. 9, 9 a .

*Hab.* PANAMA, Chiriqui<sup>1</sup> (*Mus. Madrid*).

17. **Gerridius abbreviatus.**

*Gerridius abbreviatus*, Baker, Canad. Ent. xxxix. p. 114 (1907)<sup>1</sup>.

*Hab.* NICARAGUA, San Marcos<sup>1</sup>.

Mr. Baker also records *Ischnocentrus niger* (antea, p. 155) from the same locality.

18. **Trapezoida hirsuta.**

*Trapezoida hirsuta*, Buckton, Trans. Linn. Soc., Zool. (2) ix. p. 335, t. 22. figg. 11, 11 a, b (1905)<sup>1</sup>.

*Hab.* CENTRAL AMERICA<sup>1</sup> (*Mus. Madrid*).

This new genus is said to be nearly related to *Centrotus*.

19. **Combophora besckii.**

*Combophora besckii*, Germ. Rev. Silb. iii. p. 253 (1835)<sup>1</sup>; Buckton, Mon. Membr. p. 145, t. 31. figg. 1, 1 a-c, 2, 2 a (1902)<sup>1</sup>.

*Hab.* CENTRAL AMERICA<sup>2</sup>.—BRAZIL<sup>1</sup>, Amazons<sup>2</sup>.

I feel very doubtful as to the Central-American habitat of this insect; if correct, it is the first record of the occurrence of a species of *Combophora* from our region.