

CHRYSOMPHALUS, Ashm.

1. **Chrysomphalus aonidum.**

Coccus aonidum, Linn. Syst. Nat. ed. x. p. 455 (1758).

Chrysomphalus ficus, Ashm. Amer. Entom. 1880, p. 267.


Hab. MEXICO: Tampico, Chihuahua, Matamoros, and Laguna (*Townsend*); Vera Cruz (*Cockerell*); Tlacotalpam, April 19, 1898, on a palm (*Townsend*); Orizaba, on "trueno," April 29, 1898 (*Townsend*); Vera Cruz, on a coco-nut palm, April 23, 1898 (*Townsend*). PANAMA: introduced on *Dictyosperma alba* from the Kingston Botanical Gardens (*Dolby-Tyler*).

The Linnean description of the female scale is excellent, and can apply to nothing else. This must not be confounded with *C. adonidum*.

2. **Chrysomphalus rhizophoræ.**

Chrysomphalus rhizophoræ, Ckll. Ann. & Mag. Nat. Hist., Febr. 1899, p. 169.

Hab. MEXICO: El Rio Polo, Tabasco, on mangrove (*Townsend*).



Chrysomphalus rhizophoræ, Ckll.

3. **Chrysomphalus dictyospermi.**

Aspidiotus dictyospermi, Morg. Ent. Monthly Mag. 1889, p. 352.

Hab. MEXICO: Oaxaca (*Koebele*); Cuautla (*Koebele*); ocean beach between El Faro and San Pedro, Tabasco, on rose, June 9, 1897 (*Townsend*); Minatitlan, on mango (*Townsend*). PANAMA: on *Dictyosperma alba* from the Kingston Botanical Gardens (*Dolby-Tyler*).

4. **Chrysomphalus perseæ.**

Aspidiotus perseæ, Comst. Rep. U. S. Dept. Agric. for 1880, p. 305.

Hab. MEXICO: Mazatlan (*A. de Cima*).

5. **Chrysomphalus reniformis.**

Aspidiotus reniformis, Ckll. Canad. Entom. 1897, p. 265.

Hab. MEXICO: Tehuantepec city (*Townsend*).