

6. *Chrysomphalus scutiformis*.

Aspidiotus scutiformis, Ckll. Ann. & Mag. Nat. Hist., July 1893, p. 48.

Hab. MEXICO: Soledad in Vera Cruz (*Cockerell*); Victoria (*Townsend*); Monterey (*Townsend*); Acapulco, on *Citrus* fruit from thence (*Alex. Craw*).

7. *Chrysomphalus mimosæ*.

Aspidiotus mimosæ, Comst. 2nd Rep. Dept. Entom., Cornell Univ. Exper. Sta. (1883) p. 62.

Hab. MEXICO: Tampico.

8. *Chrysomphalus bowreyi*.

Aspidiotus bowreyi, Ckll. Ent. News, 1894, p. 59.

Hab. MEXICO: Frontera (*Townsend*).

Described from Jamaica.

9. *Chrysomphalus longissimus*.

Aspidiotus longissimus, Ckll. Ann. & Mag. Nat. Hist., June 1898, p. 439.

Hab. MEXICO: Frontera (*Townsend*).

10. *Chrysomphalus albopictus*.

Aspidiotus albopictus, Ckll. Ann. & Mag. Nat. Hist., June 1898, p. 433.

Hab. MEXICO: Cuernavaca (*Townsend*); Cuautla (*Koebele*). Var. *leonis* (Twins. & Ckll.): Linares in Nuevo Leon (*Townsend*); see Journ. N. Y. Ent. Soc. 1898, p. 179.

11. *Chrysomphalus koebelei*.

Aspidiotus koebelei, Twins. & Ckll. Journ. N. Y. Ent. Soc. 1898, p. 179.

Hab. MEXICO: Oaxaca (*Koebele*).

12. *Chrysomphalus agavis*.

Aspidiotus agavis, Twins. & Ckll. Journ. N. Y. Ent. Soc. 1898, p. 178.

Hab. MEXICO: Toluca (*Koebele*); also collected by Townsend, but locality-label lost.

13. *Chrysomphalus biformis*.

Aspidiotus biformis, Ckll. Canad. Entom., May 1894, p. 131.

Hab. CENTRAL AMERICA.

Mr. A. Craw found this on an orchid from Central America.

Sect. MELANASPIS, Ckll.

14. *Chrysomphalus nigropunctatus*.

Aspidiotus nigropunctatus, Ckll. Bull. 4, Techn. Ser., Div. Ent., U. S. Dept. Agric. (1896) p. 31.

Hab. MEXICO: San Luis Potosi (*Townsend*); Apam, April 29, 1898, on "trueno" (*Townsend*); Otumba, April 29, 1898, on "trueno" (*Townsend*); Rinconada Puebla,