

**Erythrodiplax umbrata** (p. 251).

To the localities given, add :—COSTA RICA, Punta Arenas (*Bolley, colls. A. N. S., Wllmsn.* : 4 ♂, 3 ♀ b).

**Erythrodiplax ochracea** (p. 255).

To the localities given, add :—COSTA RICA, Jesus Maria [1 ♀], Punta Arenas [1 ♂, 1 ♀] (*Bolley, coll. A. N. S.*).

**Erythrodiplax connata** (p. 259).

Intermediate between **b** and **c** (p. 260).

To the localities given, add :—COSTA RICA, Juan Viñas (*Cary, U. S. N. M.* : 1 ♂).

**d** (p. 260).

To the localities given, add :—HONDURAS, Puerto Cortez [*Williamson* : 1 ♂], near San Pedro Sula [*L. A. Williamson* : 1 ♂] (*coll. Wllmsn.*).

Intermediate between **d** and **e** (p. 261).

To the localities given, add :—HONDURAS, near San Pedro Sula (*L. A. Williamson, coll. Wllmsn.* : 2 ♂).

**e** (p. 261).

To the localities given, add :—COSTA RICA, Surubres near San Mateo (*Bolley, colls. A. N. S. & Wllmsn.* : 2 ♂).

**b'** (p. 264).

To the localities given, add :—MEXICO, Baños de Santa Rosalia (*Calvert, coll. P. P. C.* : 1 ♂) in Chihuahua.

Intermediate between **b'** and **c'** (p. 264).

*Hab.* MEXICO, Baños de Santa Rosalia (*ibid.* : 1 ♂).

**c'** (p. 264).

The two following males have abdominal segments 2–7 of subequal width, 8–10 successively narrower, abd. 20–20·5, hind wing 22–24 mm.

To the localities given, add :—MEXICO, Baños de Santa Rosalia (*ibid.* : 2 ♂).

Intermediate between **c'** and **d'** (p. 266).

To the localities given, add :—MEXICO, Baños de Santa Rosalia (*ibid.* : 1 ♂), Yurecuaro in Michoacan (*ibid.* : 1 ♂).

**d'** (p. 266).

One of these males, a teneral individual, referred here on account of the extent of the colouring at the bases of the wings, has the face, including the frons, greenish-yellow.