

[SETOSA, Redt.

Setosa, Redtenbacher, op. cit. i. p. 65 (1906).

1. **Setosa versicolor**, Redt. (Tab. VIII. fig. 2.)

Setosa versicolor, Redtenbacher, op. cit. i. p. 66 (1906).

Hab. COLOMBIA, Cauca, Popayan.]

Subfam. *ANISOMORPHINÆ*.*Conspectus generum*,

- | | |
|--|-----------------------|
| 1. Femora omnia superne distincta carinata. Antennarum articulus primus depressus, intus carinatus | PARANISOMORPHA, Redt. |
| 1'. Femora antica recta, superne teretia vel obtuse carinata. Antennarum articulus primus teres | AUTOLYCA, Stål. |

PARANISOMORPHA, Redt.

Paranisomorpha, Redtenbacher, op. cit. i. p. 87 (1906).

1. **Paranisomorpha insignis**, Redt.

Paranisomorpha insignis, Redtenbacher, op. cit. i. p. 90 (1906).

Hab. COSTA RICA (*Mus. Geneva, Mus. Vienna*), Rio Sucio (*Rogers, in coll. Godman & Salvin*: 1 ♀).

AUTOLYCA, Stål.

Autolyca, Stål, Rec. Orth. iii. pp. 56, 95 (1875); Redtenbacher, op. cit. i. p. 94 (1906).

1. **Autolyca pallidicornis**, Stål.

Autolyca pallidicornis, Stål, op. cit. iii. p. 95 (1875); Redtenbacher, op. cit. i. p. 95, t. 3. figg. 18, 19.

Hab. BRITISH HONDURAS, Rio Sarstoon (*Blancaneaux, in coll. Godman & Salvin*); GUATEMALA (*Mus. Vienna, Mus. Hamburg, coll. Bolivar*), Chinautla, Capetillo, Zapote (*Champion, in coll. Godman & Salvin*), Vera Paz (*Mus. Paris*); SALVADOR (*Mus. Vienna*).—COLOMBIA.

There are three males and one female of this species in the collection before me.