

OLCYPHIDES, Griff.

Olcypoides, Griffini, Zool. Anz. xxii. p. 89 (1899) ; Redtenbacher, op. cit. i. p. 108 (1906).

1. **Olcypoides viridipes**, Rehn.

Olcypoides viridipes, Rehn, Proc. Acad. Nat. Sci. Philad. 1905, p. 798.

Hab. COSTA RICA, Caché, 1000 metres (*Mus. Acad. Nat. Sci. Philadelphia*).

This species is not included in Brunner and Redtenbacher's Monograph.

TENERELLA, Redt.

Tenerella, Redtenbacher, op. cit. i. p. 109 (1906).

1. **Tenerella tenerrima**, Redt. (Tab. VIII. fig. 4.)

Tenerella tenerrima, Redtenbacher, op. cit. i. p. 109 (1906).

Hab. PANAMA, Bugaba (*Champion, in coll. Godman & Salvin* : 1 ♀, type).—COLOMBIA (*Mus. Vienna*).

AGROSTIA, Redt.

Agrostia, Redtenbacher, op. cit. i. p. 110 (1906).

1. **Agrostia amœna**, Redt.

Agrostia amœna, Redtenbacher, op. cit. i. p. 110 (1906).

Hab. PANAMA, Volcan de Chiriqui, Bugaba (*Champion, in coll. Godman & Salvin* : 3 ♂ ♂, including the type).

BRIZOIDES, Redt.

Brizoides, Redtenbacher, op. cit. i. p. 112 (1906).

Conspectus specierum.

- 1. Elytra cum area antica alarum fusco-tincta. Alæ area postica flavescente. *B. nigricornis*, Redt.
- 1'. Elytra cum area antica alarum haud fusco-signata. Alæ area postica albida.
- 2. Antennæ virides, nigro-lineatæ. Pedes unicolores, viridi-flavescentes. *B. graminea*, Redt.
- 2'. Antennæ unicolores flavescentes. Femora annulo subapicali cum basi et apice tibiæ nigro *B. lacteipennis*, Redt.

1. **Brizoides nigricornis**, Redt. (*Briza nigricornis*, Tab. VIII. fig. 5.)

Brizoides nigricornis, Redtenbacher, op. cit. i. p. 112 (1906).

Hab. PANAMA, Chiriqui (*Mus. Vienna* : type), Bugaba (*Champion, in coll. Godman & Salvin* : 4 ♂ ♂).