

(*Mus. Oxford*), Bugaba, Chiriqui (*Champion, in coll. Godman & Salvin*), Colon (*Mus. Vienna*), Darien (*Mus. Paris*).—COLOMBIA; BRITISH GUIANA.

A long series. This species is undoubtedly very close to *P. menius*, Westw.

[EUPHASMA, Redt.

Euphasma, Redtenbacher, op. cit. i. p. 125 (1906).

1. **Euphasma bicorne**, Redt. (Tab. VIII. fig. 7.)

Euphasma bicorne, Redtenbacher, op. cit. i. p. 125 (1906).

Hab. COLOMBIA.]

PSEUDOPHASMA, Bol.

Pseudophasma, Bolivar, Act. Soc. Españ. 1896, p. 13; Redtenbacher, op. cit. i. p. 123 (1906).

1. **Pseudophasma (?) phaëton**, Rehn.

Pseudophasma phaëton, Rehn, Proc. Acad. Nat. Sci. Philad. 1904, p. 95.

Hab. COSTA RICA, San Carlos.

The systematic position of this species is very uncertain.

PLANUDES, Stål.

Planudes, Stål, Rec. Orth. iii. pp. 59, 98 (1875); Redtenbacher, op. cit. i. p. 131 (1906).

1. **Planudes crenulipes**, Rehn.

Planudes crenulipes, Rehn, Proc. Acad. Nat. Sci. Philad. 1904, p. 100.

Hab. COSTA RICA, Tucurrique.

ISAGORAS, Stål.

Isagoras, Stål, Rec. Orth. iii. p. 98 (1875); Redtenbacher, op. cit. i. p. 133 (1906).

Conspectus specierum.

1. Elytra tuberculo lato compresso, abrupte elevato, varicoso-undulato . . . *I. dentipes*, Redt.
 1'. Elytra tuberculo rotundato, haud undulato *I. plagiatus*, Redt.

1. **Isagoras dentipes**, Redt.

Isagoras dentipes, Redtenbacher, op. cit. i. p. 134 (1906).

Hab. PANAMA, Chiriqui (*Champion, in coll. Godman & Salvin, and Mus. Vienna*: 3 ♂♂, including the type).